

# Module 7: Multilevel Models for Binary Responses

## Concepts

*Fiona Steele*<sup>1</sup>  
Centre for Multilevel Modelling

### Pre-requisites

- Modules 1-3, 5, 6

## Contents

Introduction .....	1
Introduction to the Example Dataset .....	2
<b>C7.1 Two-level Random Intercept Model for Binary Responses .....</b>	<b>4</b>
C7.1.1 Generalised linear random intercept model .....	4
C7.1.2 Random intercept logit model .....	4
C7.1.3 Example: Between-state variation in voting intentions in the US .....	6
<b>C7.2 Latent Variable Representation of a Random Intercept Model for Binary Responses .....</b>	<b>11</b>
C7.2.1 Two-level random intercept threshold model .....	11
C7.2.2 Comparison of a single-level and multilevel threshold model .....	12
C7.2.3 Impact of adding a level 1 explanatory variable to a two-level model .....	14
C7.2.4 Variance partition coefficient in terms of $y^*$ .....	16
<b>C7.3 Population-Averaged and Cluster-Specific Effects .....</b>	<b>17</b>
C7.3.1 Marginal model for clustered binary data .....	17
C7.3.2 Interpretation of coefficients from random effects and marginal models .....	18
C7.3.3 Example: Comparison of marginal and random intercept models fitted to the US election data .....	22
<b>C7.4 Predicted Probabilities from a Multilevel Model .....</b>	<b>24</b>
<b>C7.5 A Two-level Random Slope Model .....</b>	<b>27</b>
C7.5.1 A random slope logit model .....	27
C7.5.2 Example: Allowing the relationship between income and voting intentions in the US to vary across states .....	28
C7.5.3 Two random coefficients: Allowing income and urban-rural differentials in voting intentions to vary across states .....	31

---

<sup>1</sup> With many thanks to Rebecca Pillinger, George Leckie, Kelvyn Jones and Harvey Goldstein for comments on earlier drafts.

<b>C7.6 Adding Level 2 Explanatory Variables: Contextual Effects .....</b>	<b>35</b>
C7.6.1 A random intercept model with a level 2 explanatory variable .....	35
C7.6.2 Cross-level interactions.....	36
<b>C7.7 Estimation of Binary Response Models.....</b>	<b>39</b>
C7.7.1 Comparison of estimation procedures .....	39
C7.7.2 Some practical guidelines on the choice between estimation procedures.....	40

## Introduction

In Module 6 we saw how multiple regression models for continuous responses can be generalised to handle binary responses. At the end of the module (C6.8), we then considered models for grouped or clustered binary data where the response variable is a proportion and the explanatory variables are defined at the group level. The application of these models was illustrated in an analysis of the proportion of voters in each state intending to vote for George Bush, including as predictors the proportion of non-white respondents in a state and the proportion who reported regular attendance at religious services.

A particular issue in the analysis of proportions is the presence of extrabinomial variation, caused by a violation of the assumption that the binary responses on which a proportion is based are independent. It was suggested in Module 6 that one way to allow for clustering (non-independence) due to omitted group-level predictors is to fit a multilevel model with group-level random effects. We pursue this approach here, but our focus is on showing how multilevel models can be applied more generally to two-level binary response data with predictors that can be defined at both level 1 and level 2.

Some examples of research questions that can be explored through multilevel models for binary responses are:

- What is the extent of between-state variation in US voting preferences (Republican vs. Democrat)? Can between-state differences in voting be explained by differences in the ethnic or religious composition of states? Do individual-level variables such as age and gender have different effects in different states?
- Does the use of dental health services (e.g. whether a person visited a dentist in the last year) vary across areas? To what extent are any differences between areas attributable to between-area differences in the provision of subsidised services or differences in the demographic and socio-economic composition of residents?

In both of the above examples, the study populations have a two-level hierarchical structure with individuals at level 1 and areas at level 2, but structures can have more than two levels and may be non-hierarchical (see Module 4). In this module, as in Module 5 for continuous responses, we consider only models for two-level hierarchical structures.

The aim of this module is to bring together multilevel models for continuous responses (Module 5) and single-level models for binary responses (Module 6). We shall see that many of the extensions to the basic multilevel model introduced in Module 5 - for example random slopes and contextual effects - apply also to binary responses. However, there are some important new issues to consider in the interpretation and estimation of multilevel binary response models.

## Introduction to the Example Dataset

We will illustrate methods for analysing binary responses using data from the 2004 National Annenberg Election Study (NAES04), a US survey designed to track the dynamics of public opinion over the 2004 presidential campaign. See <http://www.annenbergpublicpolicycenter.org> for further details of the NAES.

In this module (as in Module 6) we analyse data from the National Rolling Cross-Section of NAES04. The response variable for our analysis is based on voting intentions in the 2004 general election (variable cRC03), which was asked of respondents interviewed between 7 October 2003 and 27 January 2004. The question was worded as follows:

- *Thinking about the general election for president in November 2004, if that election were held today, would you vote for George W. Bush or the Democratic candidate?*

The response options were: Bush, Democrat, Other, Would not vote, or Depends. A small number of respondents reported that they did not know or refused to answer the question. Don't knows and refusals were excluded from the analysis, and the remaining categories were combined to obtain a binary variable coded 1 for Bush and 0 otherwise.

In Module 6 we analysed data from three states. We now extend the analysis sample to include all 49 states in the study, containing a total of 14,169 respondents.

We consider six *individual-level* explanatory variables:

- Annual household income, grouped into nine categories (1 = less than \$10k, 2 = \$10-15K, 3 = \$15-25K, 4 = \$25-35K, 5 = \$35-50K, 6 = \$50-75K, 7 = \$75-100K, 8 = \$100-150K, 9 = \$150k or more). This variable is treated as continuous in all analyses and is centred around its sample mean of 5.23
- Sex (0 = male, 1 = female)
- Age in years (mean centred)
- Type of region of residence (0 = rural, 1 = urban)
- Marital status (1 = currently married or cohabiting, 2 = widowed or divorced, 3 = not currently living with a partner and never married)

- Frequency of attendance at religious services (0 = less than weekly or never, 1 = weekly or more)

and one *state-level* explanatory variable, calculated by aggregating an individual-level variable giving the frequency of attendance at religious services:

- Proportion of respondents who attend religious services at least once a week

## C7.1 Two-level Random Intercept Model for Binary Responses

### C7.1.1 Generalised linear random intercept model

Consider a two-level structure where a total of  $n$  individuals (at level 1) are nested within  $J$  groups (at level 2) with  $n_j$  individuals in group  $j$ . Throughout this module we use ‘group’ as a general term for any level 2 unit, e.g. an area or a school. We denote by  $y_{ij}$  the response for individual  $i$  in group  $j$ , and by  $x_{ij}$  an individual-level explanatory variable. Recall from C5.2, equation (5.4), the random intercept model for continuous  $y$ :

$$y_{ij} = \beta_0 + \beta_1 x_{ij} + u_j + e_{ij} \quad (7.1)$$

where the group effects or level 2 residuals  $u_j$  and the level 1 residuals  $e_{ij}$  are assumed to be independent and to follow normal distributions with zero means:

$$u_j \sim N(0, \sigma_u^2) \text{ and } e_{ij} \sim N(0, \sigma_e^2).$$

We can also express the model in terms of the mean or *expected value* of  $y_{ij}$  for an individual in group  $j$  and with value  $x_{ij}$  on  $x$ :

$$E(y_{ij} | x_{ij}, u_j) = \beta_0 + \beta_1 x_{ij} + u_j. \quad (7.2)$$

For a binary response  $y_{ij}$ , we have  $E(y_{ij} | x_{ij}, u_j) = \pi_{ij} = \Pr(y_{ij} = 1)$  and a *generalised linear random intercept model* for the dependency of the response probability  $\pi_{ij}$  on  $x_{ij}$  is written:

$$F^{-1}(\pi_{ij}) = \beta_0 + \beta_1 x_{ij} + u_j \quad (7.3)$$

where  $F^{-1}$  (“ $F$  inverse”) is the link function, taken to be the inverse cumulative distribution function of a known distribution (see C6.3.1). In Module 6, we considered three link functions: the logit, probit and complementary log-log (clog-log) functions. Here we will focus on the logit link, with some discussion of the probit, but everything we say for the logit applies equally to the other link functions.

The key point to note about (7.3) is that, although the left hand side is a nonlinear transformation of  $\pi_{ij}$ , the right hand side takes the same form as that of (7.2) for continuous  $y$ , i.e. it is linear in terms of the parameters  $\beta_0$  and  $\beta_1$  and the level 2 residuals  $u_j$ . Therefore this simple random intercept model for binary  $y$  can be extended in the same ways that we considered in Module 5 for continuous  $y$ , including the addition of further explanatory variables defined at level 1 or 2, cross-level interactions, and random slopes (coefficients).

### C7.1.2 Random intercept logit model

In a logit model  $F^{-1}(\pi_{ij})$  is the log-odds that  $y = 1$  (see C6.3.2), so (7.3) becomes

$$\log\left(\frac{\pi_{ij}}{1-\pi_{ij}}\right) = \beta_0 + \beta_1 x_{ij} + u_j \quad (7.4)$$

where  $u_j \sim N(0, \sigma_u^2)$ .

### ***Interpretation of $\beta_0$ and $\beta_1$***

$\beta_0$  is interpreted as the *log-odds* that  $y = 1$  when  $x = 0$  and  $u = 0$  and is referred to as the *overall intercept* in the linear relationship between the log-odds and  $x$ . If we take the exponential of  $\beta_0$ ,  $\exp(\beta_0)$ , we obtain the *odds* that  $y = 1$  for  $x = 0$  and  $u = 0$ .

As in the single-level model,  $\beta_1$  is the effect of a 1-unit change in  $x$  on the log-odds that  $y = 1$ , but it is now the effect of  $x$  after adjusting for (or holding constant) the group effect  $u$ . If we are holding  $u$  constant, then we are looking at the effect of  $x$  for individuals within the same group so  $\beta_1$  is usually referred to as a *cluster-specific effect*. In C7.3 we will compare this cluster-specific effect with the effect of  $x$  averaging across groups (the *population-average effect*). These effects are equal for a multilevel continuous response model, so that in Module 5 we made no distinction between them, but they will not be equal for a generalised linear multilevel model (unless  $\sigma_u^2 = 0$ ).

As in a single-level logit model,  $\exp(\beta_1)$  can be interpreted as an odds ratio, comparing the odds that  $y = 1$  for two individuals (in the same group) with  $x$ -values spaced 1 unit apart.

### ***Interpretation of $u_j$***

While  $\beta_0$  is the overall intercept in the linear relationship between the log-odds and  $x$ , the intercept for a given group  $j$  is  $\beta_0 + u_j$  which will be higher or lower than the overall intercept depending on whether  $u_j$  is greater or less than zero. As in the continuous response case, we refer to  $u_j$  as the group (random) effect, group residual, or level 2 residual. The variance of the intercepts across groups is  $\text{var}(u_j) = \sigma_u^2$ , which is referred to as the between-group variance adjusted for  $x$ , the between-group residual variance, or simply the level 2 residual variance. (Quite often ‘residual’ is omitted and we say ‘level 2 variance’, but remember that if the model contains explanatory variables then  $\sigma_u^2$  is always the *unexplained* level 2 variance.)

We can obtain estimates of  $u_j$  that can be plotted with confidence intervals to see which groups are significantly below or above the average of zero (a caterpillar plot). These estimates are interpreted in the same way as for continuous response models (see C5.1.2 and C5.2.2); the only difference is that in a logit model they represent group effects on the log-odds scale.

In analysing multilevel data, we are often interested in the amount of variation that can be attributed to the different levels in the data structure and the extent to which variation at a given level can be explained by explanatory variables. In Module 5 (C5.1.1) we met the *variance partition coefficient* which measures the proportion of the total variance that is due to differences between groups. There is no unique

This document is only the first few pages of the full version.

To see the complete document please go to learning materials and register:

<http://www.cmm.bris.ac.uk/lemma>

**The course is completely free.** We ask for a few details about yourself for our research purposes only. We will not give any details to any other organisation unless it is with your express permission.